



Instructions:¹⁻⁶

- There are standardized instructions for each item of the BBS.
- Please refer to the protocol for standardized administration of the BBS, including instructions for each item. <http://neuropt.org/practice-resources/anpt-clinical-practice-guidelines/core-outcome-measures-cpg>

Equipment Needed:¹⁻⁶

- Stopwatch
- Standard height chair (18-20 in.) with arm rests
- Standard height chair (18-20 in.) without arm rests
- Step or stool of average height (7 $\frac{3}{4}$ - 9 in.)
- Ruler
- Slipper or shoe

Scoring:¹⁻⁶

- Items are scored on a 5 point ordinal scale.
- Points are deducted for requiring supervision, assistance and/or taking more than the allotted time to complete the task.
- The lowest category that applies should be marked.
- Please refer to the standardized administration of the BBS for item-by-item scoring. This can be found at: <http://neuropt.org/practice-resources/anpt-clinical-practice-guidelines/core-outcome-measures-cpg>

REFERENCES

1. Berg KO, Maki B, Williams JI, Holiday PJ, Wood-Dauphinee SL. Clinical and laboratory measures of postural balance in an elderly population. *Arch Phys Med Rehabil.* 1992;73(11):1073-1080.
2. Hiengkaew V, Jitaree K, Chaiyawat P. Minimal detectable changes of the Berg Balance Scale, Fugl-Meyer Assessment Scale, Timed "Up & Go" Test, gait speeds, and 2-minute walk test in individuals with chronic stroke with different degrees of ankle plantarflexor tone. *Arch Phys Med Rehabil.* 2012;93(7):1201-8.
3. Steffen T, Seney M. Test-retest reliability and minimal detectable change on balance and ambulation tests, the 36-item short-form health survey, and the unified Parkinson disease rating scale in people with parkinsonism. *Phys Ther.* 2008;88(6):733-46.
4. International Residential Code. 2012 04/17/2018]; Available from: <https://codes.iccsafe.org/public/document/toc/362/>.
5. Berg KO, Wood-Dauphinee SL, Williams JI, Maki B. Measuring balance in the elderly: validation of an instrument. *Can J Public Health.* 1992;83(Suppl 2):S7-11.
6. Berg K, Wood-Dauphinee SL, Williams JI, Gayton D. Measuring balance in the elderly: preliminary development of an instrument. *Physiother Can.* 1989;41(6):304-311.
7. Shumway-Cook A, Baldwin M, et al. Predicting the probability for falls in community-dwelling older adults. *Phys Ther.* 1997;77(8):812-819.
8. Lusardi MM, Fritz S, Middleton A, et al. Determining risk of falls in community dwelling older adults: A systematic review and meta-analysis using posttest probability. *J Geriatr Phys Ther.* 2017;40(1):1-36.
9. Stevenson TJ. Detecting change in patients with stroke using the Berg Balance Scale. *Aust J Physiother.* 2001;47(1):29-38.
10. Liaw LJ, Hsieh CL, Lo SK, Chen HM, Lee S, Lin JH. The relative and absolute reliability of two balance performance measures in chronic stroke patients. *Disabil Rehabil.* 2008;30(9):656-661.
11. Quinn L, Khalil H, Dawes H, Fritz NE, Kegelmeyer D, Kloos AD, Gillard JW, Busse M., Outcome Measures Subgroup of the European Huntington's Disease Network. Reliability and minimal detectable change of physical performance measures in individuals with pre-manifest and manifest Huntington disease. *Phys Ther.* 2013;93(7):942-956.